

In the Claims:

Please cancel claims 1-10, 15-38 and 42-59.

Please amend claims 11 and 39 as follows:

a²
11. A recombinant host cell, wherein the recombinant host cell is a *Bacillus subtilis* cell transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase.

a³
39. A method for producing hyaluronic acid, comprising the steps of:
introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain;
growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

Please add new claims 60-312.

a⁴
60. The recombinant host cell of claim 11, wherein the purified nucleic acid segment encodes a Group A hyaluronan synthase.

61. The recombinant host cell of claim 60, wherein the host cell produces hyaluronic acid.

62. The recombinant host cell of claim 11, wherein the coding region encoding enzymatically active hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

63. The recombinant host cell of claim 11, wherein the *Bacillus subtilis* cell is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

64. The recombinant host cell of claim 63, wherein the purified nucleic acid segment encodes the *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

65. The recombinant host cell of claim 63, wherein the purified nucleic acid segment comprises a nucleotide sequence in accordance with SEQ ID NO:1.

66. The recombinant host cell of claim 65, wherein the host cell produces hyaluronic acid.

67. The recombinant host cell of claim 63, wherein the purified nucleic acid segment encodes the Group A hyaluronan synthase.

68. The recombinant host cell of claim 67, wherein the host cell produces hyaluronic acid.

introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain;

introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase into the *Bacillus subtilis* strain;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and

recovering the secreted hyaluronic acid.

75. The method according to claim 74, wherein the step of recovering the hyaluronic acid comprises extracting the secreted hyaluronic acid from the medium.

ay
my. 76. The method according to claim 74, further comprising the step of purifying the extracted hyaluronic acid.

77. The method of claim 74 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the coding region encoding enzymatically active hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

78. A recombinant host cell, wherein the recombinant host cell is a *Bacillus*

subtilis cell transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

79. The recombinant host cell of claim 78, wherein the host cell produces hyaluronic acid.

80. The recombinant host cell of claim 78, wherein the coding region encoding enzymatically active Group C hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

81. The recombinant host cell of claim 78, wherein the *Bacillus subtilis* cell is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

82. A recombinant host cell, wherein the recombinant host cell is a *Bacillus subtilis* cell transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

83. The recombinant host cell of claim 82, wherein the host cell produces hyaluronic acid.

segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

90. A method for producing hyaluronic acid, comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and

recovering the secreted hyaluronic acid.

91. The method according to claim 90, wherein the step of recovering the hyaluronic acid comprises extracting the secreted hyaluronic acid from the medium.

92. The method according to claim 90, further comprising the step of purifying the extracted hyaluronic acid.

93. The method according to claim 90, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain further includes introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

94. The method according to claim 90 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the coding region encoding enzymatically active Group C hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

95. The method according to claim 90, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

ay
int.

96. The method according to claim 90, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

97. The method according to claim 90, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid

segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

98. The method according to claim 90, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

99. A method for producing hyaluronic acid, comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain;
growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

100. The method according to claim 99, wherein the step of recovering the hyaluronic acid comprises extracting the secreted hyaluronic acid from the medium.

101. The method according to claim 99, further comprising the step of purifying the extracted hyaluronic acid.

102. The method according to claim 99, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain further includes introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

103. The method according to claim 99 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

ay
cont.
104. The method according to claim 99, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

105. The method according to claim 99, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically

active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

106. The method according to claim 99, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

107. The method according to claim 99, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

108. A method for producing hyaluronic acid, comprising the steps of:
introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, wherein the purified nucleic acid segment comprises a nucleotide sequence in accordance with

FILED "3554.049"

ay
wy.

SEQ NO:1;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

109. The method according to claim 108, wherein the step of recovering the hyaluronic acid comprises extracting the secreted hyaluronic acid from the medium.

110. The method according to claim 108, further comprising the step of purifying the extracted hyaluronic acid.

111. The method according to claim 108, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain further includes introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

112. The method according to claim 108 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

113. The method according to claim 108, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

114. The method according to claim 108, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

115. The method according to claim 108, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

116. The method according to claim 108, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis*

strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

117. A recombinant host cell, wherein the recombinant host cell is a *Bacillus subtilis* cell transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase and a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

118. The recombinant host cell of claim 117, wherein the host cell produces hyaluronic acid.

ay
119. The recombinant host cell of claim 117, wherein the coding region encoding enzymatically active Group C hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

120. A recombinant host cell, wherein the recombinant host cell is a *Bacillus subtilis* cell transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 and a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

121. The recombinant host cell of claim 120, where the host cell produces hyaluronic acid.

122. The recombinant host cell of claim 120, wherein the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

123. A recombinant host cell, wherein the recombinant host cell is a *Bacillus subtilis* cell transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase and a recombinant vector comprising a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase, wherein the purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase comprises a nucleotide sequence in accordance with SEQ ID NO:1.

124. The recombinant host cell of claim 123, wherein the host cell produces hyaluronic acid.

125. The recombinant host cell of claim 123, wherein the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis*

promoter.

126. A method for producing hyaluronic acid, comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain;

introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase into the *Bacillus subtilis* strain;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and

recovering the secreted hyaluronic acid.

127. The method according to claim 126, wherein the step of recovering the hyaluronic acid comprises extracting the secreted hyaluronic acid from the medium.

128. The method according to claim 126, further comprising the step of purifying the extracted hyaluronic acid.

129. The method according to claim 126 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the coding region encoding enzymatically active Group C hyaluronan synthase of the

purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

130. The method according to claim 126, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

131. The method according to claim 126, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

132. The method according to claim 126, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

133. The method according to claim 126, wherein the step of introducing a

purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

134. A method for producing hyaluronic acid, comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain;

introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase into the *Bacillus subtilis* strain;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and

recovering the secreted hyaluronic acid.

135. The method according to claim 134, wherein the step of recovering the hyaluronic acid comprises extracting the secreted hyaluronic acid from the medium.

136. The method according to claim 134, further comprising the step of purifying the extracted hyaluronic acid.

137. The method according to claim 134 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

138. The method according to claim 134 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

139. The method according to claim 134 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

140. The method according to claim 134 wherein, in the step of introducing

a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

141. The method according to claim 134 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

142. A method for producing hyaluronic acid, comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, wherein the purified nucleic acid segment comprises a nucleotide sequence in accordance with SEQ ID NO:1;

introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase into the *Bacillus subtilis* strain;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

143. The method according to claim 142 wherein, in the step of recovering the hyaluronic acid comprises extracting the secreted hyaluronic acid from the medium.

144. The method according to claim 142, further comprising the step of purifying the extracted hyaluronic acid.

145. The method according to claim 142 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

146. The method according to claim 142 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

147. The method according to claim 142 wherein, the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

148. The method according to claim 142 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

ay
149. The method according to claim 142 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase.

150. Hyaluronic acid prepared by a process comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a

Bacillus subtilis strain;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

151. Hyaluronic acid prepared by the process according to claim 150, wherein the step of growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid is further defined as growing the *Bacillus subtilis* strain in Spizizens Minimal Media plus glucose and trace elements at about 32°C to secrete hyaluronic acid.

152. Hyaluronic acid prepared by the process according to claim 150, wherein the step of growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid is further defined as growing the *Bacillus subtilis* strain in a medium containing glucose and at least one of N-acetylglucosamine and glucosamine to secrete hyaluronic acid.

153. Hyaluronic acid prepared by the process according to claim 150, wherein the step of recovering the secreted hyaluronic acid is further defined as separating the hyaluronic acid from cells and debris by filtering and then separating the hyaluronic acid from the medium by precipitation by a precipitation agent.

154. Hyaluronic acid prepared by the process according to claim 153, wherein the step of separating the hyaluronic acid from cells and debris by filtering

further includes the addition of trichloroacetic acid which facilitates in separating cells and debris from the hyaluronic acid.

155. Hyaluronic acid prepared by the process according to claim 153, wherein the precipitation agent is an alcohol or an organic solvent.

156. Hyaluronic acid prepared by the process according to claim 155, wherein the precipitation agent is selected from the group consisting of ethanol, acetone or cetyl pyridinium chloride.

157. Hyaluronic acid prepared by the process according to claim 150, further comprising the step of purifying the extracted hyaluronic acid.

158. Hyaluronic acid prepared by the process according to claim 150, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain further includes the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

159. Hyaluronic acid prepared by the process according to claim 150 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the coding region encoding enzymatically active Group C

hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

160. Hyaluronic acid prepared by the process according to claim 150, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

161. Hyaluronic acid prepared by the process according to claim 150, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

162. Hyaluronic acid prepared by the process according to claim 150, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

163. Hyaluronic acid prepared by the process according to claim 150, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase.

164. Hyaluronic acid prepared by a process comprising the steps of:
introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain;
growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

ay
cont
165. Hyaluronic acid prepared by the process according to claim 164, wherein the step of growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid is further defined as growing the *Bacillus subtilis* strain in Spizizens Minimal Media plus glucose and trace elements at about 32°C to secrete hyaluronic acid.

166. Hyaluronic acid prepared by the process according to claim 164, wherein the step of growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid is further defined as growing the *Bacillus subtilis* strain in a medium containing glucose and at least one of N-acetylglucosamine and glucosamine to

secrete hyaluronic acid.

167. Hyaluronic acid prepared by the process according to claim 164, wherein the step of recovering the secreted hyaluronic acid is further defined as separating the hyaluronic acid from cells and debris by filtering and then separating the hyaluronic acid from the medium by precipitation by a precipitation agent.

168. Hyaluronic acid prepared by the process according to claim 167, wherein the step of separating the hyaluronic acid from cells and debris by filtering further includes the addition of trichloroacetic acid, which facilitates in separating cells and debris from the hyaluronic acid.

169. Hyaluronic acid prepared by the process according to claim 167, wherein the precipitation agent is an alcohol or an organic solvent.

170. Hyaluronic acid prepared by the process according to claim 169, wherein the precipitation agent is selected from the group consisting of ethanol, acetone or acetyl pyridinium chloride.

171. Hyaluronic acid prepared by the process according to claim 164, further comprising the step of purifying the extracted hyaluronic acid.

172. Hyaluronic acid prepared by the process according to claim 164 wherein,

in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain further includes the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

173. Hyaluronic acid prepared by the process according to claim 164 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

ay
174. Hyaluronic acid prepared by the process according to claim 164 wherein, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

175. Hyaluronic acid prepared by the process according to claim 164, wherein the step of introducing a purified nucleic acid segment having a coding region

encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

176. Hyaluronic acid prepared by the process according to claim 164, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

ay
and 177. Hyaluronic acid prepared by the process according to claim 164, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

178. Hyaluronic acid prepared by a process comprising the steps of:
introducing a purified nucleic acid segment having a coding region

encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, wherein the purified nucleic acid segment comprises a nucleotide sequence in accordance with SEQ ID NO:1;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

179. Hyaluronic acid prepared by the process according to claim 178, wherein the step of growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid is further defined as growing the *Bacillus subtilis* strain in Spizizens Minimal Media plus glucose and trace elements at about 32°C to secrete hyaluronic acid.

180. Hyaluronic acid prepared by the process according to claim 178, wherein the step of growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid is further defined as growing the *Bacillus subtilis* strain in a medium containing glucose and at least one of N-acetylglucosamine and glucosamine to secrete hyaluronic acid.

181. Hyaluronic acid prepared by the process according to claim 178, wherein the step of recovering the secreted hyaluronic acid is further defined as separating the hyaluronic acid from cells and debris by filtering and then separating the hyaluronic acid from the medium by precipitation by a precipitation agent.

182. Hyaluronic acid prepared by the process according to claim 181, wherein the step of separating the hyaluronic acid from cells and debris by filtering further includes the addition of trichloroacetic acid, which facilitates in separating cells and debris from the hyaluronic acid.

183. Hyaluronic acid prepared by the process according to claim 181, wherein the precipitation agent is an alcohol or an organic solvent.

184. Hyaluronic acid prepared by the process according to claim 183, wherein the precipitation agent is selected from the group consisting of ethanol, acetone or cetyl pyridinium chloride.

185. Hyaluronic acid prepared by the process according to claim 178, further comprising the step of purifying the extracted hyaluronic acid.

186. Hyaluronic acid prepared by the process according to claim 178, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain further includes the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active UDP-glucose dehydrogenase.

187. Hyaluronic acid prepared by the process according to claim 178 wherein, in the step of introducing a purified nucleic acid segment having a coding region

encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of the purified nucleic acid segment is under control of a *Bacillus subtilis* promoter.

188. Hyaluronic acid prepared by the process according to claim 178, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transforming a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

189. Hyaluronic acid prepared by the process according to claim 178, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transfecting a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

190. Hyaluronic acid prepared by the process according to claim 178, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase

of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as transducing a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

191. Hyaluronic acid prepared by the process according to claim 178 wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain is further defined as electroporating a *Bacillus subtilis* strain with a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2.

192. A recombinant host cell wherein the recombinant host cell is a *Bacillus subtilis* cell having an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc, the recombinant host cell further transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an enzymatically active hyaluronan synthase.

193. The recombinant host cell of claim 192, wherein the recombinant host cell further includes at least one mutated RNA polymerase promoter capable of expressing RNA polymerase in an amount greater than an endogenous RNA polymerase promoter.

194. The recombinant host cell of claim 192, wherein the recombinant host cell is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme.

195. The recombinant host cell of claim 194, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

196. The recombinant host cell of claim 194, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

197. The recombinant host cell of claim 192, wherein the recombinant host cell further includes at least one additional messenger RNA stabilizing element than is found in a native *Bacillus subtilis* cell.

ay
cx 198. The recombinant host cell of claim 192, wherein the recombinant host cell further includes at least one less messenger RNA destabilizing element than is found in a native *Bacillus subtilis* cell.

199. The recombinant host cell of claim 192, wherein the recombinant host cell further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

200. The recombinant host cell of claim 192, wherein the recombinant host cell further includes at least one mutated UDP-sugar precursor gene wherein the mutated UDP-sugar precursor gene increases a half-life of a transcribed messenger RNA.

201. The recombinant host cell of claim 192, wherein the recombinant host cell further includes at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

202. The recombinant host cell of claim 201, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased binding affinity for the ribosome binding site.

ay
203. A method for producing hyaluronic acid, comprising the steps of:
introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, wherein the *Bacillus subtilis* strain has an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc;
growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

204. The method according to claim 203, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated RNA polymerase promoter having an increased promoter activity.

205. The method according to claim 203, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain is transformed with a vector comprising a purified nucleic acid segment having a coding region encoding an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

206. The method according to claim 205, wherein the UDP-sugar precursor biosynthesis pathway is UDP-glucose dehydrogenase.

207. The method according to claim 205, wherein the UDP-sugar precursor biosynthesis pathway is UDP-glucose pyrophosphorylase.

208. The method according to claim 203, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one additional messenger RNA stabilizing element than is found in a native *Bacillus subtilis* strain.

209. The method according to claim 203, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one less messenger RNA destabilizing element than is found in a native *Bacillus subtilis* strain.

210. The method according to claim 203, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

ay.
211. The method according to claim 203, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated UDP-sugar precursor gene wherein the mutation results in an increase of a half-life of a messenger RNA transcribed from the mutated UDP-sugar precursor gene.

212. The method according to claim 203, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis*

strain further includes at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

213. The method according to claim 212, wherein the mutation to the UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a binding affinity of a ribosome for the ribosome binding site is increased.

214. A recombinant host cell wherein the recombinant host cell is a *Bacillus subtilis* cell having an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc, the recombinant host cell further being transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an enzymatically active Group C hyaluronan synthase.

ay
cont. 215. The recombinant host cell of claim 214, wherein the recombinant host cell further includes at least one mutated RNA polymerase promoter capable of expressing RNA polymerase in an amount greater than an endogenous RNA polymerase promoter.

216. The recombinant host cell of claim 214, wherein the recombinant host cell is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an UDP-sugar precursor biosynthesis pathway enzyme.

217. The recom~~inant~~ host cell of claim 216, wherein UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

218. The recombinant host cell of claim 216, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

219. The recombinant host cell of claim 214, wherein the recombinant host cell further includes at least one additional messenger RNA stabilizing element than is found in a native *Bacillus subtilis* cell.

220. The recombinant host cell of claim 214, wherein the recombinant host cell further includes at least one less messenger RNA destabilizing element than is found in a native *Bacillus subtilis* cell

ay
221. The recombinant host cell of claim 214, wherein the recombinant host cell further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

222. The recombinant host cell of claim 214, wherein the recombinant host cell further includes at least one mutated UDP-sugar precursor gene wherein the mutated UDP-sugar precursor gene increases a half-life of a transcribed messenger RNA.

223. The recombinant host cell of claim 214, wherein the recombinant host cell further includes at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

224. The recombinant host cell of claim 223, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased binding affinity for the ribosome binding site.

225. A recombinant host cell wherein the recombinant host cell is a *Bacillus subtilis* cell having an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc, the recombinant host cell further being transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an enzymatically active *Streptococcus equismilis* hyaluronan synthase of SEQ ID NO:2.

226. The recombinant host cell of claim 225, wherein the recombinant host cell further includes at least one mutated RNA polymerase promoter capable of expressing RNA polymerase in an amount greater than an endogenous RNA polymerase promoter.

227. The recombinant host cell of claim 225, wherein the recombinant host cell is transformed with a recombinant vector comprising a purified nucleic acid

segment having a coding region encoding an UDP-sugar precursor biosynthesis pathway enzyme.

228. The recombinant host cell of claim 227, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

229. The recombinant host cell of claim 227, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

230. The recombinant host cell of claim 225, wherein the recombinant host cell further includes at least one additional messenger RNA stabilizing element than is found in a native *Bacillus subtilis* cell.

ay
ent. 231. The recombinant host cell of claim 225, wherein the recombinant host cell further includes at least one less messenger RNA destabilizing element than is found in a native *Bacillus subtilis* cell.

232. The recombinant host cell of claim 225, wherein the recombinant host cell further includes at least one nucleic acid segment encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

233. The recombinant host cell of claim 225, wherein the recombinant host cell further includes at least one mutated UDP-sugar precursor gene, wherein the

mutated UDP-sugar precursor gene increases a half-life of a transcribed messenger RNA.

234. The recombinant host cell of claim 225, wherein the recombinant host cell further includes at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

235. The recombinant host cell of claim 234, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased binding affinity for the ribosome binding site.

236. A recombinant host cell wherein the recombinant host cell is a *Bacillus subtilis* cell having an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc and the recombinant host cell further being transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an enzymatically active Group C hyaluronan synthase of SEQ ID NO:1.

237. The recombinant host cell of claim 236, wherein the recombinant host cell further includes at least one mutated RNA polymerase promoter capable of expressing RNA polymerase in an amount greater than an endogenous RNA polymerase promoter.

238. The recombinant host cell of claim 236, wherein the recombinant host cell is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an UDP-sugar precursor biosynthesis pathway enzyme.

239. The recombinant host cell of claim 238, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

240. The recombinant host cell of claim 238, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

241. The recombinant host cell of claim 236, wherein the recombinant host cell further includes at least one additional messenger RNA stabilizing element than is found in a native *Bacillus subtilis* cell.

242. The recombinant host cell of claim 236, wherein the recombinant host cell further includes at least one less messenger RNA destabilizing element than is found in a native *Bacillus subtilis* cell.

243. The recombinant host cell of claim 236, wherein the recombinant host cell further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

244. The recombinant host cell of claim 236, wherein the recombinant host cell further includes at least one mutated UDP-sugar precursor gene wherein the mutated UDP-sugar precursor gene increases a half-life of a transcribed messenger RNA.

245. The recombinant host cell of claim 236, wherein the recombinant host cell further includes at least one mutated endogenous UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

246. The recombinant host cell of claim 245, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased affinity for the ribosome binding site.

247. A method for producing hyaluronic acid, comprising the steps of:
introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, wherein the *Bacillus subtilis* strain has an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc;
growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and
recovering the secreted hyaluronic acid.

248. The method according to claim 247, wherein in the step of introducing a

purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated RNA polymerase promoter having an increased promoter activity.

249. The method according to claim 247, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

250. The method according to claim 249, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

251. The method according to claim 249, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

252. The method according to claim 247, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one additional messenger RNA stabilizing element than is found in the *Bacillus subtilis* cell used to form the recombinant host cell.

253. The method according to claim 247, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one less messenger RNA destabilizing element than is found in the *Bacillus subtilis* cell used to form the recombinant host cell.

254. The method according to claim 247, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

255. The method according to claim 247, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated UDP-sugar precursor gene wherein the mutation results in an increase of a half-life of a messenger RNA transcribed from the mutated UDP-sugar precursor gene.

256. The method according to claim 247, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically

[illegible]

ay
ent

introducing a purified nucleic acid segment having a coding

equismillis hyaluronan synthase of SEQ ID NO:2 into a *Bacillus*

growing the *Bacillus subtilis* strain in a medium to secrete

recovering the secreted hyaluronic acid.

M:\GENERAL\3554 Hyalose\3554.049\PAMJP.wpd

260. The method according to claim 258, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain is transformed with a vector comprising a purified nucleic acid segment having a coding region encoding an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

261. The method according to claim 260, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

262. The method according to claim 260, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

263. The method according to claim 258, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one additional messenger RNA stabilizing element than is found in a native *Bacillus subtilis* strain.

264. The method according to claim 258, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one

less messenger RNA destabilizing element than is found in a native *Bacillus subtilis* strain.

265. The method according to claim 258, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

266. The method according to claim 258, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain comprises at least one mutated endogenous UDP-sugar precursor gene wherein the mutation results in an increase of a half-life of a messenger RNA transcribed from the mutated UDP-sugar precursor gene.

267. The method according to claim 258, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain comprises at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an

increased transcriptional efficiency.

268. The method according to claim 267, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased binding affinity for the ribosome binding site.

269. A method for producing hyaluronic acid, comprising the steps of:

introducing a purified nucleic acid segment having a coding

region encoding enzymatically active *Streptococcus*

equismillis hyaluronan synthase into a *Bacillus subtilis*

strain wherein the purified nucleic acid segment comprises a nucleotide sequence in accordance with SEQ ID NO:1, and

wherein the *Bacillus subtilis* strain has an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc;

growing the *Bacillus subtilis* strain in a medium to secrete

hyaluronic acid; and

recovering the secreted hyaluronic acid.

270. The method according to claim 269, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated RNA polymerase promoter having a greater promoter activity than an endogenous

endogenous RNA polymerase promoter.

271. The method according to claim 269, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

272. The method according to claim 271, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

273. The method according to claim 271, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

274. The method according to claim 269, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one additional messenger RNA stabilizing element than is found in a native *Bacillus subtilis* strain.

275. The method according to claim 269, wherein the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically

active *Streptococcus equismillis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one less messenger RNA destabilizing element than is found a native *Bacillus subtilis* strain.

276. The method according to claim 269, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

277. The method according to claim 269, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated endogenous UDP-sugar precursor gene wherein the mutation results in an increase of a half-life of a messenger RNA transcribed from the mutated UDP-sugar precursor gene.

278. The method according to claim 269, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equismillis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated UDP-

282. Hyaluronic acid prepared by the process according to claim 297, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

283. Hyaluronic acid prepared by the process according to claim 282, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

284. Hyaluronic acid prepared by the process according to claim 282, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

285. Hyaluronic acid prepared by the process according to claim 280, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one additional messenger RNA stabilizing element than is found in the *Bacillus subtilis* cell used to form the recombinant host cell.

286. Hyaluronic acid prepared by the process according to claim 280, wherein in the step of introducing a purified nucleic acid segment having a coding region

encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one less messenger RNA destabilizing element than is found in the *Bacillus subtilis* cell used to form the recombinant host cell.

287. Hyaluronic acid prepared by the process according to claim 280, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

288. Hyaluronic acid prepared by the process according to claim 280, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated UDP-sugar precursor gene wherein the mutation results in an increase of a half-life of a messenger RNA transcribed from the mutated UDP-sugar precursor gene.

289. Hyaluronic acid prepared by the process according to claim 280, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active Group C hyaluronan synthase into a *Bacillus*

subtilis strain, the *Bacillus subtilis* strain further includes at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

290. Hyaluronic acid prepared by the process according to claim 289, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased binding affinity for the ribosome binding site.

291. Hyaluronic acid prepared by a process comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, wherein the *Bacillus subtilis* strain has an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc;

growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and

recovering the secreted hyaluronic acid.

292. Hyaluronic acid prepared by the process according to claim 291, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated RNA polymerase promoter having a greater

promoter activity, than an endogenous RNA polymerase promoter.

293. Hyaluronic acid prepared by the process according to claim 291, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

294. Hyaluronic acid prepared by the process according to claim 293, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

295. Hyaluronic acid prepared by the process according to claim 293, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

296. Hyaluronic acid prepared by the process according to claim 291, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one additional messenger RNA stabilizing element than is

all
cont.

0507959-051301
T01E190-0506050

found in the *Bacillus subtilis* cell used to form the recombinant host cell.

297. Hyaluronic acid prepared by the process according to claim 291, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one less messenger RNA destabilizing element than is found in the *Bacillus subtilis* cell used to form the recombinant host cell.

298. Hyaluronic acid prepared by the process according to claim 291, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

299. Hyaluronic acid prepared by the process according to claim 291, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain comprises at least one mutated UDP-sugar precursor gene wherein the mutation results in an increase of a half-life of a messenger RNA transcribed from the mutated UDP-sugar precursor gene.

300. Hyaluronic acid prepared by the process according to claim 291, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase of SEQ ID NO:2 into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain comprises at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

ay
ex. 301. Hyaluronic acid prepared by the process according to claim 300, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased binding affinity for the ribosome binding site.

302. Hyaluronic acid prepared by a process for producing hyaluronic acid, comprising the steps of:

introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain wherein the purified nucleic acid segment comprises a nucleotide sequence in accordance with SEQ ID NO:1, and wherein the *Bacillus subtilis* strain has an enhanced production of at least one of UDP-GlcA and UDP-GlcNAc; growing the *Bacillus subtilis* strain in a medium to secrete hyaluronic acid; and recovering the secreted hyaluronic acid.

303. Hyaluronic acid prepared by the process according to claim 302, in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated RNA polymerase promoter having a greater promoter activity than an endogenous RNA polymerase.

304. Hyaluronic acid prepared by the process according to claim 302, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain is transformed with a recombinant vector comprising a purified nucleic acid segment having a coding region encoding an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

305. Hyaluronic acid prepared by the process according to claim 304, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose dehydrogenase.

306. Hyaluronic acid prepared by the process according to claim 304, wherein the UDP-sugar precursor biosynthesis pathway enzyme is UDP-glucose pyrophosphorylase.

307. Hyaluronic acid prepared by the process according to claim 302, wherein

in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one additional messenger RNA stabilizing element than is found in the *Bacillus subtilis* cell used to form the recombinant host cell.

308. Hyaluronic acid prepared by the process according to claim 302, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one less messenger RNA destabilizing element than is found in the *Bacillus subtilis* cell used to form the recombinant host cell.

309. Hyaluronic acid prepared by the process according to claim 302, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one nucleic acid segment having a coding region encoding a UDP-sugar precursor biosynthesis pathway enzyme having an activity greater than an endogenous UDP-sugar precursor biosynthesis pathway enzyme.

310. Hyaluronic acid prepared by the process according to claim 302, wherein in the step of introducing a purified nucleic acid segment having a coding region

encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated endogenous UDP-sugar precursor gene wherein the mutation results in an increase of a half-life of a messenger RNA transcribed from the mutated UDP-sugar precursor gene.

311. Hyaluronic acid prepared by the process according to claim 302, wherein in the step of introducing a purified nucleic acid segment having a coding region encoding enzymatically active *Streptococcus equisimilis* hyaluronan synthase into a *Bacillus subtilis* strain, the *Bacillus subtilis* strain further includes at least one mutated UDP-sugar precursor gene encoding a messenger RNA having an increased translational efficiency.

312. Hyaluronic acid prepared by the process according to claim 311, wherein the mutated UDP-sugar precursor gene occurs in a ribosome binding site in the UDP-sugar precursor gene such that a ribosome has an increased binding affinity for the ribosome binding site.